

## Real Analysis–Exam, December 11, 2002.

---

Please explain every step in detail. With a proper reference you may use anything from class and from the homework exercises.

---

**Problem 1** Consider a set  $A \subset \mathbb{R}^n$ . Show that the closure of the set obtained by forming all strictly positive convex combinations of any finite subset of  $A$  is the convex hull of  $A$ .

**Problem 2** Let  $\mathbb{X}$  be an arbitrary nonempty set and consider  $d(p, q) = \frac{1}{p+q}$  defined for  $p, q \in \mathbb{X}$ . Show that  $d$  is a metric. Determine which sets are open, closed, bounded, and compact.

**Problem 3** Let  $C \in \mathbb{R}^n$  be a convex set and let  $\delta > 0$ . Show that if  $x_1, \dots, x_k \in \mathbb{R}^n$  are such that  $D(C, x_i) < \delta$  for all  $i = 1, \dots, k$  then any convex combination  $y$  of  $x_1, \dots, x_k$  satisfies  $D(C, y) < \delta$ .

**Problem 4** Let  $\mathbb{X}, \mathbb{Y}$  be metric spaces and let  $f : \mathbb{X} \rightarrow \mathbb{Y}$  be uniformly continuous. Show that if  $\{x_n\}$  is a Cauchy sequence in  $\mathbb{X}$  then  $\{f(x_n)\}$  is Cauchy in  $\mathbb{Y}$ . Does the statement remain true if  $f$  is continuous on  $\mathbb{X}$  but not necessarily uniformly?

**Problem 5** Show that any closed and convex set  $C$  in  $\mathbb{R}^n$  equals the intersection of all closed halfspaces containing  $C$ .

**Problem 6** Let  $\mathbb{X}$  be a metric space and let  $f : \mathbb{X} \rightarrow \mathbb{X}$  be a function such that for all  $x, y \in \mathbb{X}$ ,

$$d(f(x), f(y)) < d(x, y) .$$

Give an example when  $f$  does not have any fixed point. Show that if  $\mathbb{X}$  is compact,  $f$  has exactly one fixed point. *Hint:* The function  $d(x, f(x))$  achieves its minimum on  $\mathbb{X}$ .

**Problem 7** Let  $f$  be a real-valued function defined on an open set  $E \subset \mathbb{R}^n$ , and assume that the partial derivatives  $D_1 f, \dots, D_n f$  are bounded in  $E$ . Prove that  $f$  is continuous in  $E$ .

**Problem 8** Prove that there is no continuous one-to-one and onto function  $f$  mapping the interval  $[0, 1]$  onto the square  $[0, 1] \times [0, 1]$ .