

Connectivity of sparse Bluetooth networks

N. Broutin*

L. Devroye†

G. Lugosi‡

February 12, 2015

Abstract

Consider a random geometric graph defined on n vertices uniformly distributed in the d -dimensional unit torus. Two vertices are connected if their distance is less than a “visibility radius” r_n . We consider *Bluetooth networks* that are locally sparsified random geometric graphs. Each vertex selects c of its neighbors in the random geometric graph at random and connects only to the selected points. We show that if the visibility radius is at least of the order of $n^{-(1-\delta)/d}$ for some $\delta > 0$, then a constant value of c is sufficient for the graph to be connected, with high probability. It suffices to take $c \geq \sqrt{(1+\epsilon)/\delta} + K$ for any positive ϵ where K is a constant depending on d only. On the other hand, with $c \leq \sqrt{(1-\epsilon)/\delta}$, the graph is disconnected, with high probability.

1 Introduction and results

Consider the following model of random “Bluetooth networks”. Let $\mathbf{X} = X_1, \dots, X_n$ be independent, uniformly distributed random points in $[0, 1]^d$, and denote the set of these points by $\mathbf{X} = \{X_1, \dots, X_n\}$. Given a positive number $r_n > 0$ (the so-called visibility radius), define the random geometric graph $G_n(r_n)$ with vertex set \mathbf{X} in which two vertices X_i and X_j are connected by an edge if and only if $D(X_i, X_j) \leq r_n$, where

$$D(x, y) = \left(\sum_{i=1}^d \min(|x_i - y_i|, 1 - |x_i - y_i|)^2 \right)^{1/2}$$

is the Euclidean distance on the torus.

It is well known (Penrose [11]) that, for any $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \mathbf{P}(G_n(r_n) \text{ is connected}) = \begin{cases} 0 & \text{if } r_n \leq (1 - \epsilon) \left(\frac{\log n}{nv_d} \right)^{1/d} \\ 1 & \text{if } r_n \geq (1 + \epsilon) \left(\frac{\log n}{nv_d} \right)^{1/d} \end{cases},$$

*Projet RAP, Inria Paris–Rocquencourt

†McGill University

‡ICREA and Pompeu Fabra University. GL acknowledges support by the Spanish Ministry of Science and Technology grant MTM2012-37195.

where v_d is the volume of the Euclidean unit ball in \mathbb{R}^d .

Note that in order to guarantee that a random geometric graph is connected (with high probability), the average degree in the graph needs to be at least of the order of $\log n$, which makes the graph too dense for some applications. To deal with this issue, one may sparsify the graph. This can be done in a distributed way by selecting, for each vertex X_i , randomly, and independently a subset of c_n edges adjacent to X_i , and then considering the subgraph containing the selected edges only. The selection is done without replacement and if a vertex has less than c_n neighbors in $G_n(r_n)$, then we take all of its neighbors. The obtained random graph model, coined *Bluetooth network* (or *irrigation graph*) has been introduced and studied in Crescenzi et al. [5], Dubhashi et al. [6, 7], Ferraguto et al. [9], and [12].

Formally, the random Bluetooth graph $\Gamma_n = \Gamma_n(r_n, c_n)$ is obtained as a random sub-graph of $G_n(r_n)$ as follows. For every vertex $X_i \in \mathbf{X}$, we pick randomly, without replacement, c_n edges, each adjacent to X_i in $G_n(r_n)$. (If the degree of X_i in G_n is less than c_n , all edges adjacent to X_i are kept in Γ_n .) We also denote by $\Gamma_n^+(r_n, c_n)$ the directed graph obtained by placing a directed edge from X_i to X_j whenever X_j is among the c_n selected neighbors of X_i .

We study connectivity of $\Gamma_n(r_n, c_n)$ for large values of n . A property of the graph holds *with high probability* (*whp*) when the probability that the property holds converges to one as $n \rightarrow \infty$.

When $r_n > \sqrt{d}/2$, the underlying random geometric graph $G_n(r_n)$ is the complete graph and $\Gamma_n(r_n, 2)$ becomes the “2-out” random subgraph of the complete graph studied in Fenner and Frieze [8], where it is shown that the graph is connected with high probability. Dubhashi et al. [6] extended this result by showing that when $r_n = r > 0$ is independent of n , $\Gamma_n(r, 2)$ is connected with high probability. When $r_n \rightarrow 0$ as $n \rightarrow \infty$, Crescenzi et al. [5] proved that there exist constants γ_1, γ_2 such that if $r_n \geq \gamma_1(\log n/n)^{1/d}$ and $c_n \geq \gamma_2 \log(1/r_n)$, then $\Gamma_n(r_n, c_n)$ is connected with high probability. Broutin et al. [3] proved that when r_n is just above the connectivity threshold for the underlying graph $G_n(r_n)$, that is, when $r_n \sim \gamma(\log n/n)^{1/d}$ for some sufficiently large γ , the connectivity threshold for the irrigation graphs is

$$c_n^* := \sqrt{\frac{2 \log n}{\log \log n}}.$$

More precisely, for any $\epsilon \in (0, 1)$,

$$\lim_{n \rightarrow \infty} \mathbf{P}(\Gamma_n(r_n, c_n) \text{ is connected}) = \begin{cases} 0 & \text{if } c_n \leq (1 - \epsilon)c_n^* \\ 1 & \text{if } c_n \geq (1 + \epsilon)c_n^*. \end{cases} \quad (1)$$

This result shows that the simple distributed algorithm building the irrigation graph guarantees connectivity with high probability while reducing the average degree to about $2c_n^*$, which is much less than the average degree of $\Theta(\log n)$ for the initial random geometric graph. However, the average degree still grows with n , and therefore $\Gamma_n(r_n, c_n)$ is not genuinely sparse.

MAIN RESULTS. The main result of this note is that at the price of increasing the visibility radius to $r_n \sim n^{-(1-\delta)/d}$ for some $\delta > 0$, a constant number of connections per vertex suffices to achieve connectivity with high probability. Note that with this value of r_n , the average degree of the

underlying random geometric graph is of the order of n^δ .

The lower bound of Broutin et al. [3] states that for any $\epsilon \in (0, 1)$ and $\lambda \in [1, \infty]$, if $\gamma > 0$ is a sufficiently large constant, $r_n \geq \gamma \left(\frac{\log n}{n}\right)^{1/d}$, $\frac{\log nr_n^d}{\log \log n} \rightarrow \lambda$, and

$$c_n \leq \sqrt{(1 - \epsilon) \left(\frac{\lambda}{\lambda - 1/2}\right) \frac{\log n}{\log nr_n^d}},$$

then $\Gamma_n(r_n, c_n)$ contains an isolated $(c_n + 1)$ -clique and therefore is disconnected whp.

When $r_n \sim n^{-(1-\delta)/d}$, we have $\lambda = \infty$ and therefore if $c_n \leq \lfloor (1 - \epsilon)/\sqrt{\delta} \rfloor$, then the random graph $\Gamma_n(r_n, c_n)$ is disconnected whp. This bound may seem weak since this value of c_n is just a constant, independent of n . However, we show here that this bound is essentially tight: We prove that when $r_n = \Omega(n^{-(1-\delta)/d})$, for some $\delta > 0$, then $\Gamma_n(r_n, c_n)$ is connected whp whenever c_n is larger than a certain constant, which depends on δ and d only. In fact, we show that as δ becomes small, our upper bound for the constant depends on δ as $((1 + \epsilon)/\delta)^{1/2} + O(1)$, essentially matching the lower bound of Broutin et al. [3] mentioned above (see Theorem 2 for the precise statement). To gain intuition to why one can expect a threshold value around $c \approx \delta^{-1/2}$, consider the expected number of isolated cliques of size $c + 1$ in $\Gamma_n(n^{-(1-\delta)/d}, c)$. Each vertex in $G_n(n^{-(1-\delta)/d})$ has degree concentrated around $v_d n^\delta$ (where v_d is the volume of the Euclidean ball of radius 1 in \mathbb{R}^d) and an easy argument shows that the expected number of isolated $(c + 1)$ -cliques is of the order of $n^{1-c^2\delta}$. This value is much larger than 1 when $c = ((1 - \epsilon)/\delta)^{1/2}$ but much smaller than 1 when $c = ((1 + \epsilon)/\delta)^{1/2}$. As it often happens in other random graph models, the last obstacles for connectivity are the smallest possible isolated components and as soon as isolated $(c + 1)$ -cliques disappear, the graph becomes connected. However, our proof of connectivity follows a different path, showing the existence of a path between any two vertices.

Theorem 1. *Let $\delta \in (0, 1)$, $\gamma > 0$, and $\epsilon \in (0, 1)$ be fixed. Suppose that $r_n = \gamma_n n^{-(1-\delta)/d}$ where $\gamma_n/\gamma \rightarrow 1$ as $n \rightarrow \infty$. There exists a constant c , depending on δ and d only, such that the random Bluetooth graph $\Gamma_n(r_n, c)$ is connected whp. For $x \in (0, 1)$ set*

$$f(x) := \sqrt{(1 + x^2 + 8\sqrt{x} + \epsilon)/(x - 2x^2 \log_2(1/x))}.$$

One may take $c = k_1 + k_2 + k_3 + 1$, where

$$k_1 = \begin{cases} \lceil f(\delta) \rceil & \text{if } \delta \in (0, 1/5) \\ \lceil f(1/5) \rceil & \text{if } \delta \in (1/5, 1) \end{cases}, \quad k_2 = \left\lceil \frac{8(1 + \epsilon)v_d(2\sqrt{d})^d}{(1 - \epsilon)} \right\rceil, \quad k_3 = \lceil \sqrt{4(1 + \epsilon)v_d/\alpha_d} \rceil$$

where v_d is the volume of the Euclidean ball of radius 1 in \mathbb{R}^d and $\alpha_d = (1 - \epsilon)/(2(2\sqrt{d})^d)$.

A straightforward combination of the lower bound and Theorem 1 implies the following:

Theorem 2. *Let $\bar{c}(\delta)$ denote the smallest integer c for which*

$$\lim_{n \rightarrow \infty} \mathbf{P}(\Gamma_n(r_n, c) \text{ is connected}) = 1$$

and let $\underline{c}(\delta)$ denote the largest integer c for which

$$\lim_{n \rightarrow \infty} \mathbf{P}(\Gamma_n(r_n, c) \text{ is disconnected}) = 1$$

when $r_n = \gamma_n n^{-(1-\delta)/d}$ with $\gamma_n \sim \gamma$ as $n \rightarrow \infty$. Then, for any $\epsilon > 0$, we have, for all δ small enough,

$$(1 - \epsilon)\delta^{-1/2} \leq \underline{c}(\delta) \leq \bar{c}(\delta) \leq (1 + \epsilon)\delta^{-1/2}.$$

Interestingly, the threshold is essentially independent of the value of γ and the dimension d . This phenomenon was also observed in (1).

REMARKS AND OPEN QUESTIONS. Before we conclude this section, we mention a few questions that might be worth investigating. Theorem 2 above finds the correct asymptotics for the thresholds $\underline{c}(\delta), \bar{c}(\delta)$ when the visibility radius is $r_n \sim \gamma n^{-(1-\delta)/d}$ for small δ . One may phrase a related question as follows: given a fixed a constant c , (a budget, in some sense) can one find a threshold function $r_n^* = r_n^*(c)$ for connectivity? More precisely, r_n^* should be such that, for any $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \mathbf{P}(\Gamma_n(r_n, c) \text{ is connected}) = \begin{cases} 0 & \text{if } r_n < (1 - \epsilon)r_n^* \\ 1 & \text{if } r_n > (1 + \epsilon)r_n^*. \end{cases}$$

In some sense, Theorem 2 gives the asymptotics of the threshold function for $c \rightarrow \infty$, but one would like to know the threshold for fixed values of the budget c . As it was proved in [3] in the case that $r_n \sim \gamma(n^{-1} \log n)^{1/d}$, the main obstacles to connectivity should be isolated $(c + 1)$ -cliques. One could also try to prove that this is indeed the case at a finer level: around the threshold, the number of isolated $(c + 1)$ -cliques should be asymptotically Poisson distributed. Thus, one expects that the probability that $\Gamma_n(r_n, c)$ is connected should have asymptotics similar to that of classical random graphs [1, Theorem 7.3] or random geometric graphs [11, Theorem 13.11], where the isolated vertices are the main obstacle.

Finally, we mention that elsewhere (Broutin et al. [4]) we investigate the birth of the giant component of $\Gamma_n(r_n, c)$.

2 Proof sketch

In the course of the proof, we condition on the location of the points and assume that they are sufficiently regularly distributed. The probability that this happens is estimated in the following lemma. Let $N(A) = \sum_{i=1}^n \mathbf{1}_{\{X_i \in A\}}$ denote the number of points falling in a set $A \subset [0, 1]^d$ and let λ denote the Lebesgue measure. Denote by $B_{x,r} = \{y \in [0, 1]^d : D(x, y) < r\}$ the open ball centered at $x \in [0, 1]^d$ and by $C_{x,r} = \{y : \forall i = 1, \dots, d, \min(|x_i - y_i|, 1 - |x_i - y_i|) \leq r/2\}$ the cube of side length r centered at $x \in [0, 1]^d$.

Lemma 1. *Suppose $r_n \geq \gamma n^{-(1-\delta)/d}$ for some $\delta \in (0, 1)$ and $\gamma > 0$. Let $\epsilon > 0$ and denote by F*

the event that for all $x \in [0, 1]^d$,

$$\frac{N(B_{x,r_n})}{n\lambda(B_{x,r_n})} \in (1 - \epsilon, 1 + \epsilon) \quad \text{and} \quad \frac{N(C_{x,r_n/(2\sqrt{d})})}{n\lambda(C_{x,r_n/(2\sqrt{d})})} \in (1 - \epsilon, 1 + \epsilon).$$

Then there exists a constant $\theta = \theta(\delta, \epsilon) > 0$ and a positive integer $n_0 = n_0(\delta, \epsilon)$ such that for all $n > n_0$, $\mathbf{P}(F) \geq 1 - \exp(-\theta nr_n^d)$.

Proof. The lemma may be proved by standard arguments. It follows, for example, from inequalities for uniform deviations of empirical measures over VC classes such as Theorem 7 in [2]. \square

In the rest of the proof, we fix $\mathbf{X} = \{X_1, \dots, X_n\}$, and assume that event F holds. Thus, all randomness originates from the random choices of the c neighbors of every vertex. We denote by \mathbf{P}_c the probability with respect to the random choice of the neighbors only (i.e., conditional given the set \mathbf{X}). It suffices to show that if F holds, then with high probability (with respect to \mathbf{P}_c), $\Gamma(r_n, c)$ is connected.

SKETCH OF THE PROOF. The general strategy is to prove that, with high probability, from any two points X_i, X_j , one can find a path that connects X_i to X_j . To do this, most of the work consists in proving structural properties of the connected component containing a fixed point X_1 . We study the random graph by dividing the c neighbors into four disjoint groups of sizes k_1, k_2, k_3 , and 1, thus obtaining four independent sets of edges added in four different phases. The sketch of the proof of Theorem 1 is as follows. We rely on a discretization of the unit cube $[0, 1]^d$ into congruent cubes of side length $1/\lceil 2\sqrt{d}/r_n \rceil$.

- SEARCHING FOR A DENSE CUBE I. (Section 3). In the first phase we start from an arbitrary vertex, say, X_1 , and using only k_1 choices of each vertex, consider the set of the vertices of \mathbf{X} which may be reached using paths of at most $\ell \approx \delta^2 \log_2 n$ edges. We show that if this growth process succeeds, there exists a cube in the grid partition that contains a connected component of size at least $n^{\min\{\delta^2, 1/25\}}/3$.
- SEARCHING FOR A DENSE CUBE II. (Section 4). In the second phase we show that by adding k_2 new connections to each vertex in the component obtained after the first step, at least one of the grid cells has a positive fraction of its points in a single connected component.
- PROPAGATING THE DENSITY. (Section 5). Once we have a cell containing a constant proportion of points belonging to the same connected component, it is rather easy to propagate this positive density (of a single connected component) to all other cells of the grid by using k_3 new connections per vertex.
- CONNECTIVITY IS UNAVOIDABLE. (Section 6). The previous phases guarantee that from a single vertex X_1 all three phases succeed with probability $1 - o(1/n)$. So, with high probability, the connected components of every single vertex X_i , $1 \leq i \leq n$, reach in every corner of the space. Then, it is easy to show that with just one additional connection per

vertex, any two such components very likely connect, proving that the entire graph is, in fact, connected.

3 First growth process: searching for a dense cube I

Divide $[0, 1]^d$ into a grid of congruent cubes of side length $1/\lceil 2\sqrt{d}/r_n \rceil$. We first prove that a constant number of edges per vertex suffices to guarantee that there exists a cell that contains at least a polynomial number of points of \mathbf{X} that may be reached from X_1 .

Lemma 2. *Suppose $\mathbf{X} = \{X_1, \dots, X_n\}$ are such that the event F defined in Lemma 1 occurs. Let k_1 and r_n be defined as in Theorem 1. With probability at least $1 - o(1/n)$, the connected component of $\Gamma(r_n, k_1)$ containing X_1 is such that there exists a cell in the grid partition of $[0, 1]^d$ into congruent cubes of side length $1/\lceil 2\sqrt{d}/r_n \rceil$ that contains at least $n^{\min\{\delta^2, 1/25\}/3}$ vertices of the component.*

Proof. Let $\ell < n$ be a positive integer specified below. Consider A , the set of vertices of \mathbf{X} that can be reached from X_1 using a directed path of length at most ℓ in $\Gamma_n^+(r_n, k_1)$. Note that $|A| \leq 1 + k_1 + k_1^2 + \dots + k_1^\ell$. We first show a lower bound on the size of this connected component:

$$\mathbf{P}(|A| \leq k_1 2^{\ell-1}) = o(1/n). \quad (2)$$

To see this, the key property is that with high probability, the number of new points added in the second generation is at least $2k_1$. First, the k_1 points of the first generation are distinct for they are sampled without replacement. For the second generation, imagine k_1^2 bins, k_1 for each of the k_1 vertices of the first generation, into which we place the points chosen by these vertices. For these k_1^2 bins to contain only j different points that are also different from the points of the first generation, there must exist $k_1^2 - j$ bins that contain only points of the j remaining bins or points from the first generation. There are $\binom{k_1^2}{j}$ ways of choosing these $k_1^2 - j$ bins and for each such bin, the probability that it contains a point either from the remaining j bins or from points of the first generation is at most $(1 + k_1 + k_1^2)/(nr_n^d v_d(1 - \epsilon) - 1 - k_1 - k_1^2)$ (since, on the event F , each ball of radius r_n contains at least $nr_n^d v_d(1 - \epsilon)$ points). Thus, the probability that the number of distinct points in the second generation that are distinct and do not belong to the first generation is less than $2k_1$ is at most

$$\sum_{j=0}^{2k_1} \binom{k_1^2}{j} \left(\frac{1 + k_1 + k_1^2}{nr_n^d v_d(1 - \epsilon) - 1 - k_1 - k_1^2} \right)^{k_1^2 - j}.$$

Similarly, assuming that there are at least $2k_1$ points in the second generation, the probability that the number of selected neighbors in the third generation not selected before is less than $4k_1$ is at most

$$\sum_{j=0}^{4k_1} \binom{2k_1^2}{j} \left(\frac{1 + k_1 + k_1^2 + k_1^3}{nr_n^d v_d(1 - \epsilon) - 1 - k_1 - k_1^2 - k_1^3} \right)^{2k_1^2 - j}.$$

We may continue in this fashion for $\ell - 1$ steps, in each step doubling the number of neighbors with high probability. The probability that the ℓ -th generation has less than $2^{\ell-1}k_1$ vertices is at most

$$\begin{aligned} & \sum_{i=1}^{\ell-1} \sum_{j=0}^{2^{i-1}k_1} \binom{2^{i-1}k_1^2}{j} \left(\frac{1 + k_1 + \dots + k_1^i}{nr_n^d v_d(1-\epsilon) - (1 + k_1 + \dots + k_1^i)} \right)^{2^{i-1}k_1^2 - j} \\ & \leq \sum_{i=1}^{\ell-1} 2^{i-1}k_1 2^{2^{i-1}k_1^2} \left(\frac{k_1^\ell}{nr_n^d v_d(1-\epsilon) - k_1^\ell} \right)^{2^{i-1}k_1^2 - 2^i k_1}. \end{aligned} \quad (3)$$

Now, we choose $\ell = \ell(\delta) = \lfloor \min\{\delta^2, 1/25\} \log_2 n \rfloor$, and distinguish two cases depending on the value of δ .

(i) Suppose first that $\delta \in (0, 1/5]$. In this case, we aim at obtaining a value for k_1 that matches the lower bound. Then, in this range,

$$\log_2 k_1 \leq \log_2 \left(\sqrt{\frac{1 + \delta^2 + 8\sqrt{\delta} + \epsilon}{\delta - 2\delta^2 \log_2(1/\delta)}} + 1 \right) \leq 2 \log_2(1/\delta),$$

for any $\epsilon \in (0, 1)$ and $\delta \in (0, 1/5]$. Thus, we have $k_1^\ell \leq n^{\delta^2 \log_2 k_1} \leq n^{2\delta^2 \log_2(1/\delta)}$. It follows that the right-hand side in (3) above is at most

$$\begin{aligned} & \sum_{i=1}^{\ell-1} 2^{i-1}k_1 \left(2^{1/(1-2/k_1)} \right)^{2^{i-1}k_1^2(1-2/k_1)} \left(\frac{n^{2\delta^2 \log_2(1/\delta)}}{\gamma_n^d n^\delta v_d(1-\epsilon) - n^{2\delta^2 \log_2(1/\delta)}} \right)^{2^{i-1}k_1^2(1-2/k_1)} \\ & \leq \sum_{i=1}^{\ell-1} 2^{i-1}k_1 \left(\frac{2^{1/(1-2/k_1)}}{\gamma_n^d v_d(1-2\epsilon)} n^{-\delta+2\delta^2 \log_2(1/\delta)} \right)^{2^{i-1}k_1^2(1-2/k_1)} \\ & \leq \ell 2^\ell k_1 \kappa_d^{k_1^2} n^{-k_1^2(1-2/k_1)(\delta-2\delta^2 \log_2(1/\delta))} \\ & \leq k_1 \kappa_d^{k_1^2} \cdot n^{\delta^2 \log_2(n)} \cdot n^{-k_1^2(1-2/k_1)(\delta-2\delta^2 \log_2(1/\delta))} \end{aligned}$$

for n sufficiently large, where $\kappa_d = 8/(\gamma^d v_d(1-2\epsilon))$. Now, by our choice of k_1 , we have

$$\begin{aligned} & \delta^2 - k_1^2(1-2/k_1)(\delta-2\delta^2 \log_2(1/\delta)) \\ & \leq \delta^2 - (1 + \delta^2 + 8\sqrt{\delta} + \epsilon) + 2\sqrt{(1 + \delta^2 + 8\sqrt{\delta} + \epsilon)(\delta-2\delta^2 \log_2(1/\delta))} \\ & \leq -1 - 8\sqrt{\delta} - \epsilon + 2\sqrt{12\delta} \\ & \leq -1 - \epsilon, \end{aligned}$$

hence the probability in (3) above is $o(1/n)$ and the bound in (2) is proved for $\delta \in (0, 1/5]$.

(ii) Suppose next that $\delta \in (1/5, 1)$. In this case, the bound follows trivially from the fact that the

case $\delta = 1/5$ is covered by case (i). Indeed, we have

$$\begin{aligned} & \sum_{i=1}^{\ell-1} 2^{i-1} k_1 2^{2^{i-1} k_1^2} \left(\frac{k_1^\ell}{nr_n^d v_d (1-\epsilon) - k_1^\ell} \right)^{2^{i-1} k_1^2 - 2^i k_1} \\ & \leq \sum_{i=1}^{\ell-1} 2^{i-1} k_1 \left(\frac{4k_1^{\ell(1/5)}}{nr_n^d v_d (1-\epsilon) - k_1^{\ell(1/5)}} \right)^{2^{i-1} k_1^2 (1-2/k_1)}. \end{aligned}$$

It follows that, in this range also, the probability in (3) is $o(1/n)$ so that the bound (2) is proved for $\delta \in (1/5, 1)$.

Thus, we have shown that if n is sufficiently large, then with probability at least $1 - o(1/n)$, A contains at least $k_1 2^{\ell-1} \geq n^{\min\{\delta^2, 1/25\}/2}$ vertices, all within distance $\ell r_n \leq \min\{\delta^2, 1/25\} \log_2 n \cdot \gamma_n n^{-(1-\delta)/d}$ from X_1 . In particular, the points of A all fall in grid cells at most $\lceil r_n \ell \rceil / (r_n / (2\sqrt{d})) + 1 \leq 1 + 2\ell\sqrt{d}$ away from the cell containing X_1 . Thus, all these vertices fall in a cube of at most $(3 + 4\ell\sqrt{d})^d$ cells. This implies that there must exist a cell with at least

$$\frac{n^{\min\{\delta^2, 1/25\}/2}}{(3 + 4\ell\sqrt{d})^d} \geq n^{\min\{\delta^2, 1/25\}/3}$$

vertices for n large enough. □

4 Second growth process: searching for a dense cube II

We now show that we can leverage Lemma 2 and obtain, still using a constant number of extra edges per vertex, a cell that contains a positive density of points of the connected component containing X_1 .

Lemma 3. *Suppose event F occurs. Let k_1, k_2 , and r_n be defined as in Theorem 1. With probability at least $1 - o(1/n)$, the connected component of $\Gamma_n(r_n, k_1 + k_2)$ containing X_1 is such that there exists a cell in the grid partition of $[0, 1]^d$ into congruent cubes of side length $1/\lceil 2\sqrt{d}/r_n \rceil$ that contains at least $\alpha_d nr_n^d$ vertices of the component, where $\alpha_d = (1 - \epsilon)/(2(2\sqrt{d})^d)$.*

By Lemma 2 we see that, with probability at least $1 - o(1/n)$, after k_1 connections per vertex, there exists a grid cell that contains at least $n^{\min\{\delta^2, 1/25\}/3}$ vertices of the connected component containing X_1 . Next we show that with

$$k_2 = \left\lceil \frac{8(1 + \epsilon)v_d(2\sqrt{d})^d}{(1 - \epsilon)} \right\rceil$$

new connections, the same cell contains a constant times nr_n^d vertices in the same connected component (i.e., a linear fraction of all points in the cell).

Consider the cell that contains the largest number of vertices in the connected component containing X_1 after the first k_1 connections, and let N_0 denote the number of such connected

vertices in this cell. Then we have seen in Section 3 that

$$\mathbf{P}(N_0 < n^{\min\{\delta^2, 1/25\}/3}) = o(1/n)$$

with respect to the random choices of the first k_1 connections. Suppose that the event $N_0 \geq n^{\min\{\delta^2, 1/25\}/3}$ holds.

Now add k_2 fresh connections to each of these N_0 points, resulting in $N_1 \leq k_2 N_0$ vertices in the same grid cell that haven't been in the connected component of X_1 so far. If the number N_1 of new vertices in the cell added to the component is less than $2n^{\min\{\delta^2, 1/25\}/3}$, we declare failure, otherwise continue by adding k_2 new connections to these N_1 points. (Note that since these N_1 vertices did not belong to the component of X_1 before the first step, we have not discovered any of their connections and we may use k_2 new connections per vertex.) In this step we add $N_2 \leq k_2 N_1$ new vertices in the same grid cell. If $N_2 < 4n^{\min\{\delta^2, 1/25\}/3}$, we declare failure, otherwise continue. We repeat adding k_2 connections to all newly discovered vertices until the number of connected vertices in the grid cell $N_0 + N_1 + \dots + N_i$ reaches $\alpha_d n r_n^d$ where $\alpha_d = (1 - \epsilon)/(2(2\sqrt{d})^d)$ or else for L steps, requiring in every step $i = 1, \dots, L$ that the number of newly discovered vertices in the same cell be at least $2^i n^{\min\{\delta^2, 1/25\}/3}$. Here L is chosen such that

$$2^L n^{\min\{\delta^2, 1/25\}/3} \leq \alpha_d n r_n^d < 2^{L+1} n^{\min\{\delta^2, 1/25\}/3}.$$

To estimate the probability that the process described above fails, observe first that at step i , the (conditional) probability that a vertex selects a *new* neighbor in the same cell is at least

$$\frac{2\alpha_d n r_n^d - \sum_{j=0}^{i-1} N_j}{(1 + \epsilon) n r_n^d v_d} \geq p_d \stackrel{\text{def}}{=} \frac{\alpha_d}{(1 + \epsilon) v_d}$$

since on the event F , every grid cell has at least $(1 - \epsilon) n r_n^d / (2\sqrt{d})^d = 2\alpha_d n r_n^d$ points and the vertex can reach at most $(1 + \epsilon) n r_n^d v_d$ points. In the inequality we used the fact that the number of vertices discovered until step i during the process $\sum_{j=0}^{i-1} N_j \leq (1 - \epsilon) n r_n^d / (2(2\sqrt{d})^d)$ otherwise the process stops with success before step i .

Thus, after $i - 1$ successful steps, the expected number of newly discovered vertices at stage i is at least

$$\mathbf{E}_i N_i \geq N_{i-1} k_2 p_d \geq 2^{i-1} n^{\min\{\delta^2, 1/25\}/3} k_2 p_d \geq 2^{i+1} n^{\min\{\delta^2, 1/25\}/3}$$

by the definition of k_2 , where \mathbf{E}_i denotes conditional expectation given the first growth process and the first $i - 1$ steps of the second process. Given that the process has not failed up to step $i - 1$,

the conditional probability that it fails at step i is thus

$$\begin{aligned} \mathbf{P}_i(N_i < 2^i n^{\min\{\delta^2, 1/25\}/3}) &\leq \mathbf{P}_i(N_i < \mathbf{E}_i N_i - 2^i n^{\min\{\delta^2, 1/25\}/3}) \\ &\leq \exp\left(\frac{-k_2 N_{i-1} p_d^2}{8}\right) \\ &\leq \exp\left(\frac{-k_2 2^{i-1} n^{\min\{\delta^2, 1/25\}/3} p_d^2}{8}\right), \end{aligned}$$

where we used the fact that conditionally, N_i has a hypergeometric distribution whose moment generating function is dominated by that of the corresponding binomial distribution $B(k_2 N_{i-1}, p_d)$ (see Hoeffding [10]) and we used simple Chernoff bounding for the binomial distribution.

Thus, the probability that the process ever fails is bounded by

$$\begin{aligned} \sum_{i=1}^L \exp\left(\frac{-k_2 2^{i-1} n^{\min\{\delta^2, 1/25\}/3} p_d^2}{8}\right) &\leq L \exp\left(\frac{-k_2 n^{\min\{\delta^2, 1/25\}/3} p_d^2}{8}\right) \\ &\leq \exp\left(-n^{\min\{\delta^2, 1/25\}/4}\right) \end{aligned}$$

for all sufficiently large n .

5 Third growth process: propagating the density

In this third step we show that, by adding a few more connections, the connected component containing X_1 contains a linear fraction of the points in every cell of the grid partition of $[0, 1]^d$ into cubes of side length $1/\lceil 2\sqrt{d}/r_n \rceil$. In order to do so, we start from the cell containing $\alpha_d n r_n^d$ vertices in the connected component of $\Gamma_n(r_n, k_1 + k_2)$ containing X_1 , and “grow” the component cell-by-cell until every cell has the required number of vertices in the same connected component. We show that a constant number of additional connections per vertex suffices.

Lemma 4. *Suppose event F defined in Lemma 1 occurs. Let k_1, k_2, k_3 , and r_n be defined as in Theorem 1. With probability at least $1 - o(1/n)$, the connected component of $\Gamma_n(r_n, k_1 + k_2 + k_3)$ containing X_1 is such that every cell in the grid partition of $[0, 1]^d$ into congruent cubes of side length $1/\lceil 2\sqrt{d}/r_n \rceil$ contains at least $2\alpha_d n r_n^d / (3k_3)$ vertices of the component.*

Proof. Suppose the event described in Lemma 3 holds so that there exists a cell with $\alpha_d n r_n^d$ vertices in the connected component of X_1 . Label this cell by 1. Next label all the cells from 1 to $\lceil 2\sqrt{d}/r_n \rceil^d$ in such a way that cells i and $i + 1$ are adjacent (i.e., they share a $d - 1$ -dimensional face), for all $i = 1, \dots, \lceil 2\sqrt{d}/r_n \rceil^d - 1$. Note that the size of the cells was chosen such that for any x in cell i the ball $B(x, r)$ contains entirely cell $i + 1$. In particular, every point in cell $i + 1$ is a potential neighbor of a point in cell i .

Let $k_3 = \lceil \sqrt{4(1 + \epsilon)v_d/\alpha_d} \rceil$ and consider the following process. Select, arbitrarily,

$$M := \left\lfloor \frac{2\alpha_d n r_n^d}{3k_3} \right\rfloor$$

of the already connected vertices in cell 1 and select, one-by-one, k_3 new neighbors for each until M newly connected vertices have been found in cell 2. Declare failure if the number of newly connected vertices in cell 2 is less than M after revealing all possible k_3M new connections. Otherwise continue and select k_3 new neighbors of the new vertices in cell 2. Visit all cells in a sequential fashion, always stopping the selection when M new vertices are connected in the next cell. If the process succeeds, then X_1 is connected to at least M points in every cell, which is a positive proportion.

To analyze the probability of failure, note that for each new connection, the probability of discovering a new vertex in cell $i + 1$ by a vertex in cell i is at least

$$\frac{2\alpha_d n r_n^d - k_3 M}{(1 + \epsilon) v_d n r_n^d}$$

since (assuming the event F) there are at least $2\alpha_d n r_n^d$ vertices in cell $i + 1$, at most $(1 + \epsilon) v_d n r_n^d$ within radius r_n of the point whose neighbors we select, and we discard at most $k_3 M$ points cell $i + 1$ that may have been already be chosen by other vertices in cell i .

Thus, the expected number of newly discovered vertices in cell $i + 1$, conditionally on the fact that the process has not failed earlier, is at least

$$k_3 \alpha_d n r_n^d \cdot \frac{2\alpha_d n r_n^d - k_3 M}{(1 + \epsilon) v_d n r_n^d} \geq k_3 \alpha_d n r_n^d \frac{\alpha_d}{(1 + \epsilon) v_d} \geq 2M ,$$

for all n large enough, where the last inequality follows from our choice of k_3 . Thus, the probability of failure in step i is, by a similar argument as in the proof of Lemma 3, bounded by $e^{-\eta_d n r_n^d}$, where

$$\eta_d = \frac{\alpha_d^3}{12k_3(1 + \epsilon)^2 v_d^2} .$$

Hence, the probability that the process ever fails in any step is $o(1/n)$ and this completes the proof. \square

6 Final step: proof of Theorem 1

With a giant component densely populating every cell of the grid, it is now easy to show that with at most one extra connection, the entire graph becomes connected, with high probability. The argument is as follows.

Start by growing the component of X_1 as in the growth processes described above. Then, with $k_1 + k_2 + k_3$ connections per vertex, the component has the property described in Lemma 4, with probability $1 - o(1/n)$. Now consider points X_2, X_3, \dots, X_n , one-by-one. If X_2 does not belong to the connected component of X_1 , then grow the same process starting from X_2 until the component of X_2 hits the one of X_1 . If the two components do not connect, then this new component also satisfies the property of Lemma 4, with probability $1 - o(1/n)$. (Note that until the two components meet, all connections are new in the sense that they have not been revealed in the first process, and therefore these events hold independently.) Now we may continue, taking

every X_i , $i = 3, \dots, n$. If X_i does not belong to any of the previously grown components, then we grow a new component starting from X_i until it hits one of the previous ones, or else until the component has at least $2\alpha_d n r_n^d / (3k_3)$ vertices in every cell of the grid. By the union bound, with probability $1 - o(1)$, every vertex is contained in such a giant component.

If there is more than one connected component at the end of the process, then we add one more connection to every vertex not belonging to the component of X_1 . Note that each such component has at least $(1 - \epsilon)n / (3k_3)$ vertices and for every vertex, the probability that it hits the component of X_1 is at least $\eta := 2\alpha_d / (3k_3 v_d (1 + \epsilon))$: indeed, on the event F , every ball of radius r_n^d contains at most $n r_n^d v_d (1 + \epsilon)$ points, while it fully contains a cell with $2\alpha_d n r_n^d / (3k_3)$ points of the connected component of X_1 . As a consequence the probability that all new connections miss the component of X_1 is at most $\eta^{(1 - \epsilon)n / (3k_3)}$. Thus, by the union bound, with high probability, all components connect to the first one and the entire graph becomes connected. This concludes the proof of Theorem 1.

Acknowledgement

Part of this work has been conducted while attending a workshop at the Banff International Research Station for the workshop on Models of Sparse Graphs and Network Algorithms.

References

- [1] B. Bollobás. *Random Graphs*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2001.
- [2] O. Bousquet, S. Boucheron, and G. Lugosi. Introduction to statistical learning theory. In O. Bousquet, U.v. Luxburg, and G. Rätsch, editors, *Advanced Lectures in Machine Learning*, pages 169–207. Springer, 2004.
- [3] N. Broutin, L. Devroye, N. Fraiman, and G. Lugosi. Connectivity threshold for Bluetooth graphs. *Random Structures & Algorithms*, 2013. To appear.
- [4] N. Broutin, L. Devroye, and G. Lugosi. Almost optimal sparsification of random geometric graphs. 2014. Manuscript.
- [5] P. Crescenzi, C. Nocentini, A. Pietracaprina, and G. Pucci. On the connectivity of Bluetooth-based ad hoc networks. *Concurrency and Computation: Practice and Experience*, 21(7): 875–887, 2009.
- [6] D. Dubhashi, C. Johansson, O. Häggström, A. Panconesi, and M. Sozio. Irrigating ad hoc networks in constant time. In *Proceedings of the Seventeenth Annual ACM Symposium on Parallelism in Algorithms and Architectures*, pages 106–115. ACM, 2005.
- [7] D. Dubhashi, O. Häggström, G. Mambrini, A. Panconesi, and C. Petrioli. Blue pleiades, a new solution for device discovery and scatternet formation in multi-hop Bluetooth networks. *Wireless Networks*, 13(1):107–125, 2007.

- [8] T.I. Fenner and A.M. Frieze. On the connectivity of random m -orientable graphs and digraphs. *Combinatorica*, 2:347–359, 1982.
- [9] F. Ferraguto, G. Mambrini, A. Panconesi, and C. Petrioli. A new approach to device discovery and scatternet formation in Bluetooth networks. In *Proceedings of the 18th International Parallel and Distributed Processing Symposium*, 2004.
- [10] W. Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58:13–30, 1963.
- [11] M. Penrose. *Random Geometric Graphs*, volume 5 of *Oxford Studies in Probability*. Oxford University Press, Oxford, 2003.
- [12] A. Pettarin, A. Pietracaprina, and G. Pucci. On the expansion and diameter of Bluetooth-like topologies. *Algorithms-ESA 2009*, pages 528–539, 2009.